

**Package insert**

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use **AFLURIA® QUADRIVALENT** safely and effectively. See full prescribing information for **AFLURIA QUADRIVALENT**.

**AFLURIA QUADRIVALENT, Influenza Vaccine**  
**Suspension for Intramuscular Injection**

**2018-2019 Formula**

**Initial U.S. Approval (AFLURIA QUADRIVALENT): 2016**

-----**RECENT MAJOR CHANGES**-----

Indications and Usage (1) 07/2017  
Dosage and Administration (2) 07/2017

-----**INDICATIONS AND USAGE**-----

- **AFLURIA QUADRIVALENT** is an inactivated influenza vaccine indicated for active immunization against influenza disease caused by influenza A subtype viruses and type B viruses contained in the vaccine. (1)
- **AFLURIA QUADRIVALENT** is approved for use in persons 6 months of age and older. (1)

-----**DOSAGE AND ADMINISTRATION**-----

**For intramuscular injection only, by needle and syringe (6 months and older) or by PharmaJet®Stratis® Needle-Free Injection System (18 through 64 years).** (2)

Age	Dose	Schedule
6 months through 35 months	One or two doses <sup>a</sup> , 0.25 mL each	If 2 doses, administer at least 1 month apart
36 months through 8 years	One or two doses <sup>a</sup> , 0.5 mL each	If 2 doses, administer at least 1 month apart
9 years and older	One dose, 0.5 mL	Not Applicable

<sup>a</sup>1 or 2 doses depends on vaccination history as per Advisory Committee on Immunization Practices annual recommendations on prevention and control of influenza with vaccines. (2)

-----**DOSAGE FORMS AND STRENGTHS**-----

**AFLURIA QUADRIVALENT** is a suspension for injection supplied in three presentations:

- 0.25 mL pre-filled syringe (single dose) (3, 11)
- 0.5 mL pre-filled syringe (single dose) (3, 11)
- 5 mL multi-dose vial (ten 0.5 mL doses) (3, 11)

-----**CONTRAINDICATIONS**-----

- Severe allergic reaction (e.g., anaphylaxis) to any component of the vaccine including egg protein, or to a previous dose of any influenza vaccine. (4, 11)

-----**WARNINGS AND PRECAUTIONS**-----

- If Guillain-Barré Syndrome (GBS) has occurred within 6 weeks of previous influenza vaccination, the decision to give **AFLURIA QUADRIVALENT** should be based on careful consideration of the potential benefits and risks. (5.1)
- Appropriate medical treatment and supervision must be available to manage possible anaphylactic reactions following administration of the vaccine. (5.2)

-----**ADVERSE REACTIONS**-----

**AFLURIA QUADRIVALENT** administered by needle and syringe:

- In adults 18 through 64 years, the most commonly reported injection-site adverse reaction was pain (≥ 40%). The most common systemic adverse events were myalgia and headache (≥ 20%). (6.1)
- In adults 65 years of age and older, the most commonly reported injection-site adverse reaction was pain (≥ 20%). The most common systemic adverse event was myalgia (≥ 10%). (6.1)
- In children 5 through 8 years, the most commonly reported injection-site adverse reactions were pain (≥ 50%), redness and swelling (≥ 10%). The most common systemic adverse event was headache (≥ 10%). (6.1)
- In children 9 through 17 years, the most commonly reported injection-site adverse reactions were pain (≥ 50%), redness and swelling (≥ 10%). The most common systemic adverse events were headache, myalgia, and malaise and fatigue (≥ 10%). (6.1)
- In children 6 months through 35 months of age, the most commonly reported injection-site reactions were pain and redness (≥ 20%). The most common systemic adverse events were irritability (≥ 30%), diarrhea and loss of appetite (≥ 20%). (6.1)
- In children 36 through 59 months of age, the most commonly reported injection site reactions were pain (≥ 30%) and redness (≥ 20%). The most commonly reported systemic adverse events were malaise and fatigue, and diarrhea (≥ 10%). (6.1)

**AFLURIA** (trivalent formulation) administered by the PharmaJet Stratis Needle-Free Injection System:

- In adults 18 through 64 years of age, the most commonly reported injection-site adverse reactions were tenderness (≥ 80%), swelling, pain, redness (≥ 60%), itching (≥ 20%) and bruising (≥ 10%). The most common systemic adverse events were myalgia, malaise (≥ 30%), and headache (≥ 20%). (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Seqirus USA Inc. at 1-855-358-8966 or VAERS at 1-800-822-7967 or [www.vaers.hhs.gov](http://www.vaers.hhs.gov).**

-----**USE IN SPECIFIC POPULATIONS**-----

- The safety and effectiveness of **AFLURIA QUADRIVALENT** in persons less than 6 months of age have not been established. (8.4)
- Antibody responses were lower in geriatric subjects than in younger adults. (8.5)
- Pregnancy: There is a pregnancy exposure registry that monitors outcomes in women exposed to **AFLURIA QUADRIVALENT** during pregnancy. Enroll in the pregnancy registry by calling 1-855-358-8966 or sending an email to [us.medicalinformation@seqirus.com](mailto:us.medicalinformation@seqirus.com). (8.1)

See 17 for **PATIENT COUNSELING INFORMATION**.

**Revised: 06/2018**

## Package insert

---

### FULL PRESCRIBING INFORMATION: CONTENTS\*

- 1 INDICATIONS AND USAGE**
  - 2 DOSAGE AND ADMINISTRATION**
  - 3 DOSAGE FORMS AND STRENGTHS**
  - 4 CONTRAINDICATIONS**
  - 5 WARNINGS AND PRECAUTIONS**
    - 5.1 Guillain-Barré Syndrome
    - 5.2 Preventing and Managing Allergic Reactions
    - 5.3 Altered Immunocompetence
    - 5.4 Limitations of Vaccine Effectiveness
  - 6 ADVERSE REACTIONS**
    - 6.1 Clinical Trials Experience
    - 6.2 Postmarketing Experience
  - 7 DRUG INTERACTIONS**
  - 8 USE IN SPECIFIC POPULATIONS**
    - 8.1 Pregnancy
    - 8.2 Lactation
    - 8.4 Pediatric Use
    - 8.5 Geriatric Use
  - 11 DESCRIPTION**
  - 12 CLINICAL PHARMACOLOGY**
    - 12.1 Mechanism of Action
  - 13 NONCLINICAL TOXICOLOGY**
    - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
  - 14 CLINICAL STUDIES**
    - 14.1 Efficacy Against Laboratory-Confirmed Influenza
    - 14.2 Immunogenicity of AFLURIA QUADRIVALENT in Adults and Older Adults Administered by Needle and Syringe
    - 14.3 Immunogenicity of AFLURIA (trivalent formulation) Administered by PharmaJet Stratis Needle-Free Injection System
    - 14.4 Immunogenicity of AFLURIA QUADRIVALENT in Children 5 through 17 Years Administered by Needle and Syringe
    - 14.5 Immunogenicity of AFLURIA QUADRIVALENT in Children 6 Months through 59 Months Administered by Needle and Syringe
  - 15 REFERENCES**
  - 16 HOW SUPPLIED/STORAGE AND HANDLING**
    - 16.1 How Supplied
    - 16.2 Storage and Handling
  - 17 PATIENT COUNSELING INFORMATION**
- \* Sections or subsections omitted from the full prescribing information are not listed.

**Package insert**

---

**1 FULL PRESCRIBING INFORMATION****1 INDICATIONS AND USAGE**

AFLURIA<sup>®</sup> QUADRIVALENT is an inactivated influenza vaccine indicated for active immunization against influenza disease caused by influenza A subtype viruses and type B viruses contained in the vaccine.

AFLURIA QUADRIVALENT is approved for use in persons 6 months of age and older.

**2 DOSAGE AND ADMINISTRATION****For intramuscular (IM) use only.**

- By needle and syringe (6 months of age and older)
- By PharmaJet<sup>®</sup> Stratis<sup>®</sup> Needle-Free Injection System (18 through 64 years of age)

The dose and schedule for AFLURIA QUADRIVALENT are presented in Table 1.

**Table 1: AFLURIA QUADRIVALENT Dosage and Schedule**

Age	Dose	Schedule
6 months through 35 months	One or two doses <sup>a</sup> , 0.25 mL each	If 2 doses, administer at least 1 month apart
36 months through 8 years	One or two doses <sup>a</sup> , 0.5 mL each	If 2 doses, administer at least 1 month apart
9 years and older	One dose, 0.5mL	Not Applicable

<sup>a</sup>1 or 2 doses depends on vaccination history as per Advisory Committee on Immunization Practices annual recommendations on prevention and control of influenza with vaccines.

Immediately before use, shake thoroughly and inspect visually. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever suspension and container permit. If either of these conditions exists, the vaccine should not be administered.

When using the single-dose pre-filled syringe, shake the syringe thoroughly and administer the dose immediately.

When using the multi-dose vial, shake the vial thoroughly before withdrawing each dose, and administer the dose immediately.

- Needle and Syringe: Draw up the exact dose using a separate sterile needle and syringe for each individual patient. It is recommended that small syringes (0.5 mL or 1 mL) be used to minimize any product loss.
- PharmaJet Stratis Needle-Free Injection System: For instructions on withdrawal of a 0.5 mL dose and use of the PharmaJet Stratis Needle-Free Injection System, refer to the Instructions For Use for the PharmaJet Stratis Needle-Free Injection System.

The preferred sites for intramuscular injection are the anterolateral aspect of the thigh in

**Package insert**

---

32 infants 6 months through 11 months of age, the anterolateral aspect of the thigh (or the deltoid  
33 muscle of the upper arm if muscle mass is adequate) in persons 12 months through 35 months  
34 of age, or the deltoid muscle of the upper arm in persons  $\geq 36$  months of age.

**3 DOSAGE FORMS AND STRENGTHS**

35 AFLURIA QUADRIVALENT is a sterile suspension for intramuscular injection (*see*  
36 *Description [11]*).

37  
38 AFLURIA QUADRIVALENT is supplied in three presentations:

- 39 • 0.25 mL pre-filled syringe (single dose, for persons 6 months through 35 months of  
40 age)
- 41 • 0.5 mL pre-filled syringe (single dose, for persons 36 months of age and older).
- 42 • 5 mL multi-dose vial ( for persons 6 months of age and older).

**4 CONTRAINDICATIONS**

43  
44 AFLURIA QUADRIVALENT is contraindicated in individuals with known severe allergic  
45 reactions (e.g., anaphylaxis) to any component of the vaccine including egg protein, or to a  
46 previous dose of any influenza vaccine (*see Description [11]*).

**5 WARNINGS AND PRECAUTIONS****5.1 Guillain-Barré Syndrome**

48 If Guillain-Barré Syndrome (GBS) has occurred within 6 weeks of previous influenza  
49 vaccination, the decision to give AFLURIA QUADRIVALENT should be based on careful  
50 consideration of the potential benefits and risks.  
51

52 The 1976 swine influenza vaccine was associated with an increased frequency of GBS.  
53 Evidence for a causal relation of GBS with subsequent vaccines prepared from other influenza  
54 viruses is unclear. If influenza vaccine does pose a risk, it is probably slightly more than one  
55 additional case per 1 million persons vaccinated.

**5.2 Preventing and Managing Allergic Reactions**

56 Appropriate medical treatment and supervision must be available to manage possible  
57 anaphylactic reactions following administration of the vaccine.  
58

**5.3 Altered Immunocompetence**

59 If AFLURIA QUADRIVALENT is administered to immunocompromised persons, including  
60 those receiving immunosuppressive therapy, the immune response may be diminished.  
61

**5.4 Limitations of Vaccine Effectiveness**

62 Vaccination with AFLURIA QUADRIVALENT may not protect all individuals.  
63

**Package insert**

---

**6 ADVERSE REACTIONS**

64  
65 In adults 18 through 64 years of age, the most commonly reported injection-site adverse  
66 reaction observed in clinical studies with AFLURIA QUADRIVALENT administered by  
67 needle and syringe was pain ( $\geq 40\%$ ). The most common systemic adverse events observed  
68 were myalgia and headache ( $\geq 20\%$ ).

69 In adults 65 years of age and older, the most commonly reported injection-site adverse reaction  
70 observed in clinical studies with AFLURIA QUADRIVALENT administered by needle and  
71 syringe was pain ( $\geq 20\%$ ). The most common systemic adverse event observed was myalgia  
72 ( $\geq 10\%$ ).

73 The safety experience with AFLURIA (trivalent formulation) is relevant to AFLURIA  
74 QUADRIVALENT because both vaccines are manufactured using the same process and have  
75 overlapping compositions (see [Description \[11\]](#)).

76 In adults 18 through 64 years of age, the most commonly reported injection-site adverse  
77 reactions observed in a clinical study with AFLURIA (trivalent formulation) using the  
78 PharmaJet Stratis Needle-Free Injection System were tenderness ( $\geq 80\%$ ), swelling, pain,  
79 redness ( $\geq 60\%$ ), itching ( $\geq 20\%$ ) and bruising ( $\geq 10\%$ ). The most common systemic adverse  
80 events were myalgia, malaise ( $\geq 30\%$ ) and headache ( $\geq 20\%$ ).

81 In children 5 through 8 years, the most commonly reported injection-site adverse reactions  
82 when AFLURIA QUADRIVALENT was administered by needle and syringe were pain ( $\geq$   
83  $50\%$ ) and redness and swelling ( $\geq 10\%$ ). The most common systemic adverse event was  
84 headache ( $\geq 10\%$ ).

85 In children 9 through 17 years, the most commonly reported injection-site adverse reactions  
86 when AFLURIA QUADRIVALENT was administered by needle and syringe were pain ( $\geq$   
87  $50\%$ ) and redness and swelling ( $\geq 10\%$ ). The most common systemic adverse events were  
88 headache, myalgia, and malaise and fatigue ( $\geq 10\%$ ).

89 In children 6 months through 35 months of age, the most frequently reported injection site  
90 reactions in the clinical study with AFLURIA QUADRIVALENT administered by needle and  
91 syringe were pain and redness ( $\geq 20\%$ ). The most common systemic adverse events were  
92 irritability ( $\geq 30\%$ ), diarrhea and loss of appetite ( $\geq 20\%$ ).

93 In children 36 through 59 months of age, the most commonly reported injection site reactions  
94 were pain ( $\geq 30\%$ ) and redness ( $\geq 20\%$ ). The most commonly reported systemic adverse  
95 events were malaise and fatigue, and diarrhea ( $\geq 10\%$ ).

96

**6.1 Clinical Trials Experience**

97  
98 Because clinical studies are conducted under widely varying conditions, adverse reaction rates  
99 observed in the clinical studies of a vaccine cannot be directly compared to rates in the clinical  
100 studies of another vaccine and may not reflect the rates observed in clinical practice.

**Package insert**

---

101 *Adults*

102 Clinical safety data for AFLURIA QUADRIVALENT in adults have been collected in one  
103 clinical trial, Study 1, a randomized, double-blind, active-controlled trial conducted in the U.S.  
104 in 3449 subjects ages 18 years and older. Subjects in the safety population received one dose  
105 of either AFLURIA QUADRIVALENT (N=1721) or one of two formulations of comparator  
106 trivalent influenza vaccine (AFLURIA, TIV-1 N=864 or TIV-2 N=864) each containing an  
107 influenza type B virus that corresponded to one of the two B viruses in AFLURIA  
108 QUADRIVALENT (a type B virus of the Yamagata lineage or a type B virus of the Victoria  
109 lineage), respectively. The mean age of the population was 58 years, 57% were female, and  
110 racial groups consisted of 82% White, 16% Black, and 2% other; 5% of subjects were  
111 Hispanic/Latino. The age sub-groups were 18 through 64 years and 65 years and older with  
112 mean ages of 43 years and 73 years, respectively. In this study, AFLURIA QUADRIVALENT  
113 and comparator trivalent influenza vaccines were administered by needle and syringe (*see*  
114 *Clinical Studies [14]*).

115 Local (injection-site) adverse reactions and systemic adverse events were solicited for 7 days  
116 post-vaccination (Table 2). Injection site cellulitis, cellulitis-like reactions (defined as  
117 concurrent Grade 3 pain, redness, and swelling/lump), and Grade 3 swelling/lump were  
118 monitored for 28 days post-vaccination. Unsolicited adverse events were collected for 28 days  
119 post-vaccination. Serious adverse events (SAEs), including deaths, were collected for 180  
120 days post-vaccination.

Package insert

121 **Table 2: Proportion of Subjects Per Age Cohort with Any Solicited Local Adverse**  
122 **Reactions or Systemic Adverse Events within 7 Days after Administration of**  
123 **AFLURIA QUADRIVALENT or Trivalent Influenza Vaccine (Study 1)<sup>a</sup>**

	Percentage (%) <sup>b</sup> of Subjects in each Age Cohort Reporting an Event											
	Subjects 18 through 64 years						Subjects ≥ 65 years					
	AFLURIA Quadrivalent N= 854 <sup>c</sup>		TIV-1 N= 428 <sup>c</sup>		TIV-2 N= 430 <sup>c</sup>		AFLURIA Quadrivalent N= 867 <sup>c</sup>		TIV-1 N= 436 <sup>c</sup>		TIV-2 N= 434 <sup>c</sup>	
	Any	Gr 3	Any	Gr 3	Any	Gr 3	Any	Gr 3	Any	Gr 3	Any	Gr 3
<b>Local Adverse Reactions <sup>d</sup></b>												
Pain	47.9	0.7	43.7	1.4	50.7	1.2	24.6	0.1	22.7	0	21.0	0.2
Swelling/Lump	3.7	0.1	2.3	0	3.5	0.2	3.2	0.5	1.8	0	1.6	0
Redness	2.9	0	2.8	0	2.8	0	4.2	0.3	2.1	0	2.5	0.2
<b>Systemic Adverse Events <sup>e</sup></b>												
Myalgia (muscle ache)	25.5	1.9	23.4	1.4	24.2	1.2	12.7	0.3	14.0	0.7	12.2	0.5
Headache	21.7	1.7	15.2	0.9	19.1	1.2	8.4	0	7.1	0.2	7.8	0.7
Malaise	8.9	0.7	9.1	0	9.3	0.7	4.4	0.5	5.0	0.2	5.1	0.2
Nausea	6.9	0.6	7.7	0.5	6.3	1.2	1.6	0	1.8	0	2.1	0.2
Chills	4.8	0.6	4.4	0.2	4.7	0.5	2.0	0	2.1	0.5	1.4	0.2
Vomiting	1.5	0.4	0.9	0	2.3	0.7	0.5	0.1	0	0	0.7	0.2
Fever	1.1	0.4	0.9	0	0.5	0	0.2	0	0.9	0	0.5	0.2

124 Abbreviations: Gr 3, Grade 3.

125 <sup>a</sup> NCT02214225

126 <sup>b</sup> Proportion of subjects reporting each solicited local adverse reaction or systemic adverse event by study vaccine group based  
127 on the number of subjects contributing any follow up safety information for at least one data value of an individual  
128 sign/symptom.

129 <sup>c</sup> N = number of subjects in the Safety Population for each study vaccine group.

130 <sup>d</sup> Local adverse reactions: Grade 3 pain is that which prevents daily activity; Swelling/Lump and redness: any = ≥ 20mm  
131 diameter, Grade 3 = ≥ 100mm diameter.

132 <sup>e</sup> Systemic adverse events: Fever: any = ≥ 100.4°F (Oral), Grade 3 = ≥ 102.2°F (Oral); Grade 3 for all other adverse events is  
133 that which prevents daily activity.

134 In the 28 days following vaccination, no subject experienced cellulitis or a cellulitis-like  
135 reaction. All Grade 3 swelling/lump reactions began within 7 days of vaccination and are  
136 included in Table 2.

137 In the 28 days following vaccination, 20.5%, 20.1%, and 20.7% of adults 18 through 64 years  
138 and 20.3%, 24.1%, and 20.0% of adults ≥ 65 years who received AFLURIA  
139 QUADRIVALENT, TIV-1, and TIV-2, respectively, reported unsolicited adverse events.  
140 Rates of individual events were similar between treatment groups, and most events were mild  
141 to moderate in severity.

142 In the 180 days following vaccination, 2.3%, 1.6%, and 1.5% of all subjects who received

**Package insert**

---

143 AFLURIA QUADRIVALENT, TIV-1, and TIV-2, respectively, experienced SAEs, including  
144 six deaths, five in the AFLURIA QUADRIVALENT group and one in the TIV-2 group. The  
145 majority of SAEs occurred after Study Day 28 and in subjects  $\geq 65$  years of age who had co-  
146 morbid illnesses. No SAEs or deaths appeared related to the study vaccines.

147 Safety information has also been collected in a clinical study of AFLURIA (trivalent  
148 formulation) administered using the PharmaJet Stratis Needle-Free Injection System (Study 2).  
149 Study 2 included 1,247 subjects for safety analysis, ages 18 through 64 years, randomized to  
150 receive AFLURIA by either the PharmaJet Stratis Needle-Free Injection System (624 subjects)  
151 or needle and syringe (623 subjects). No deaths or vaccine-related serious adverse events were  
152 reported in Study 2. Local (injection-site) adverse reactions and systemic adverse events were  
153 solicited for 7 days post-vaccination (Table 3).



Package insert

154 **Table 3: Proportion of Subjects 18 through 64 Years of Age with Solicited Local Adverse**  
 155 **Reactions or Systemic Adverse Events within 7 Days after Administration of**  
 156 **AFLURIA (trivalent formulation) by PharmaJet Stratis Needle-Free Injection**  
 157 **System or Needle and Syringe (Study 2)<sup>a</sup>**

	Percentage <sup>b</sup> of Subjects Reporting Event			
	Subjects 18 through 64 years			
	AFLURIA (trivalent formulation)			
	PharmaJet Stratis Needle-Free Injection System N=540-616 <sup>c</sup>		Needle and Syringe N=599-606 <sup>c</sup>	
	Any	Grade 3	Any	Grade 3
<b>Local Adverse Reactions <sup>d</sup></b>				
Tenderness	89.4	2.1	77.9	1.0
Swelling	64.8	1.7	19.7	0.2
Pain	64.4	0.8	49.3	0.7
Redness	60.1	1.3	19.2	0.3
Itching <sup>f</sup>	28.0	0.0	9.5	0.2
Bruising	17.6	0.2	5.3	0.0
<b>Systemic Adverse Events <sup>e</sup></b>				
Myalgia	36.4	0.8	35.5	1.0
Malaise	31.2	0.7	28.4	0.5
Headache	24.7	1.3	22.1	1.3
Chills	7.0	0.2	7.2	0.2
Nausea	6.6	0.2	6.5	0.0
Vomiting	1.3	0.0	1.8	0.2
Fever	0.3	0.0	0.3	0.0

158 <sup>a</sup> NCT01688921

159 <sup>b</sup> Proportion of subjects reporting each local adverse reaction or systemic adverse event by treatment group based on the  
 160 number of subjects contributing at least one data value for an individual sign/symptom (individual event denominators).

161 <sup>c</sup> N = number of subjects in the Safety Population for each treatment group. Denominators for the PharmaJet Stratis Needle-  
 162 Free Injection System group were: N=540 for itching and N=605-616 for all other parameters. Denominators for the needle  
 163 and syringe group were: N=527 for itching and N=599-606 for all other parameters.

164 <sup>d</sup> Local adverse reactions: Grade 3 is pain, tenderness or itching that prevents daily activity; Swelling, redness or bruising: any  
 165 = ≥ 25mm diameter, Grade 3 = > 100mm diameter.

166 <sup>e</sup> Systemic adverse events: Fever: any = ≥ 100.4°F (Oral), Grade 3 = ≥ 102.2°F (Oral); Grade 3 for all other adverse events is  
 167 that which prevents daily activity.

168 <sup>f</sup> A total of 155 subjects (approximately randomly distributed between PharmaJet Stratis Needle-Free Injection System and  
 169 needle and syringe groups) received Diary Cards without itching listed as a solicited symptom.

170 In adults 18 through 64 years who received AFLURIA (trivalent formulation) administered by  
 171 PharmaJet Stratis Needle-Free Injection System, commonly reported unsolicited adverse  
 172 events were headache (4.2%), injection site hematoma (1.8%), injection site erythema (1.1%),  
 173 myalgia (1.0%) and nausea (1.0%).

**Package insert**

---

174 ***Children 5 Years Through 17 Years of Age***

175 Clinical safety data for AFLURIA QUADRIVALENT in older children and adolescents have  
176 been collected in one clinical trial, Study 3, a randomized, observer-blinded, comparator-  
177 controlled trial conducted in the U.S. in 2278 subjects aged 5 through 17 years. Subjects were  
178 stratified into one of two age cohorts of 5 through 8 years or 9 through 17 years (51.2% and  
179 48.8% of the study population, respectively). The mean age of the population was 9.5 years,  
180 52.1% were male, and racial groups consisted of 73.3% White, 20.7% Black, 0.8% Asian,  
181 0.3% American Indian/Native American, and 0.7% Native Hawaiian/Pacific Islander; 23.8% of  
182 subjects were Hispanic/Latino. The mean ages of subjects 5 through 8 years and 9 through 17  
183 years were 6.7 years and 12.5 years, respectively. Subjects in the safety population (N=2252)  
184 received either AFLURIA QUADRIVALENT (N=1692) or a U.S.-licensed comparator  
185 quadrivalent influenza vaccine (N=560). Study subjects were scheduled to receive either a  
186 single vaccination or two vaccinations 28 days apart based on their previous vaccination  
187 history. In this study, AFLURIA QUADRIVALENT and comparator vaccine were  
188 administered by needle and syringe (see *Clinical Studies [14]*).

189 Local (injection site) adverse reactions and systemic adverse events were solicited for 7 days  
190 post-vaccination. Cellulitis-like reactions (defined as concurrent Grade 3 pain, redness, and  
191 swelling/lump) at the injection site were monitored for 28 days post-vaccination. Subjects  
192 were instructed to report and return to clinic within 24 hours in the event of a cellulitis-like  
193 reaction. Unsolicited adverse events were collected for 28 days post-vaccination. All solicited  
194 local adverse reactions and systemic adverse events following any vaccination (first or second  
195 dose) are presented in Table 4.

Package insert

196 **Table 4: Proportion of Subjects Per Age Cohort with Any Solicited Local Adverse**  
197 **Reactions or Systemic Adverse Events within 7 Days after Administration of**  
198 **AFLURIA QUADRIVALENT or Comparator (Study 3)<sup>a</sup>**

	Percentage (%) <sup>b</sup> of Subjects in each Age Cohort Reporting an Event							
	Subjects 5 through 8 years				Subjects 9 through 17 years			
	AFLURIA Quadrivalent N= 828-829 <sup>c</sup>		Comparator N= 273-274 <sup>c</sup>		AFLURIA Quadrivalent N= 790-792 <sup>c</sup>		Comparator N= 261 <sup>c</sup>	
	Any	Gr 3	Any	Gr 3	Any	Gr 3	Any	Gr 3
<b>Local Adverse Reactions <sup>d</sup></b>								
Pain	51.3	0.8	49.6	0.7	51.5	0.3	45.2	0.4
Redness	19.4	3.5	18.6	1.8	14.8	1.9	16.1	1.9
Swelling/Lump	15.3	3.4	12.4	2.2	12.2	2.0	10.7	1.9
<b>Systemic Adverse Events <sup>e</sup></b>								
Headache	12.3	0.1	10.6	0.4	18.8	0.4	14.6	0.4
Myalgia	9.8	0.1	11.3	0.4	16.7	0.3	11.1	0.4
Malaise and Fatigue	8.8	0.4	5.8	0	10.0	0.4	7.7	0
Nausea	7.1	0.1	8.4	0	7.7	0	8.0	0
Diarrhea	5.2	0	3.6	0	5.4	0	4.2	0
Fever	4.5	1.2	3.6	0.7	2.1	0.5	0.8	0
Vomiting	2.4	0.2	4.4	0	1.8	0	2.3	0

199 Abbreviations: Gr 3, Grade 3 (severe); Comparator, Comparator quadrivalent influenza vaccine [Fluarix<sup>®</sup> Quadrivalent  
200 (GlaxoSmithKline Biologicals)]

201 <sup>a</sup> NCT02545543

202 <sup>b</sup> Percent (%) is derived from the number of subjects that reported the event divided by the number of subjects in the Solicited  
203 Safety Population with non-missing data for each age cohort, treatment group, and each solicited parameter.

204 <sup>c</sup> N = number of subjects in the Solicited Safety Population (subjects who were vaccinated and provided any solicited safety  
205 data) for each study vaccine group.

206 <sup>d</sup> Local adverse reactions: Grade 3 pain is that which prevents daily activity; swelling/lump and redness: any = > 0mm  
207 diameter, Grade 3 = > 30mm diameter.

208 <sup>e</sup> Systemic adverse events: Fever: any = ≥ 100.4°F (Oral), Grade 3 = ≥ 102.2°F (Oral); Grade 3 for all other adverse events is  
209 that which prevents daily activity or requires significant medical intervention.  
210

211 In subjects 5 through 8 years of age, all solicited local adverse reactions and systemic adverse  
212 events were reported at lower frequencies after the second vaccination than after the first  
213 vaccination with AFLURIA QUADRIVALENT with the exception of vomiting (which  
214 occurred at the same rate of 2.2% after each vaccination).

215 One subject, 8 years of age, experienced a cellulitis-like reaction at the injection site after  
216 vaccination with AFLURIA QUADRIVALENT.

217 The most commonly reported unsolicited adverse events in the 28 days following the first or  
218 second dose of AFLURIA QUADRIVALENT in subjects 5 through 8 years of age were cough  
219 (2.4%), pyrexia (1.8%), rhinorrhea (1.2%), and headache (1.0%), and were similar to the  
220 comparator.

**Package insert**

---

221 For subjects ages 9 through 17 years who received AFLURIA QUADRIVALENT, the most  
222 commonly reported unsolicited adverse events in the 28 days following vaccination were  
223 oropharyngeal pain (1.6%), cough (1.3%), and upper respiratory tract infection (1.0%), and  
224 were similar to the comparator.

225 No deaths were reported in Study 3. In the 180 days following vaccinations, AFLURIA  
226 QUADRIVALENT and comparator vaccine recipients experienced similar rates of serious  
227 adverse events (SAEs). None of the SAEs appeared related to the study vaccines except for  
228 one case of influenza B infection (considered a vaccine failure) in an AFLURIA  
229 QUADRIVALENT recipient.

**230 *Children 6 Months Through 59 Months of Age***

231 Clinical safety data for AFLURIA QUADRIVALENT in infants and young children have been  
232 collected in one clinical trial, Study 4, a randomized, observer-blind, comparator-controlled  
233 trial conducted in the U.S. in 2247 subjects aged 6 through 59 months. Subjects were stratified  
234 into one of two age cohorts of 6 through 35 months or 36 through 59 months (41.6% and  
235 58.4% of the study population, respectively). The mean age of the population was 36.6  
236 months, 51.6% were male, and racial groups consisted of 71.0% White, 21.5% Black, 1.1%  
237 Asian, 0.7% Native Hawaiian/Pacific Islander, and 0.3% American Indian/Native American;  
238 26.4% of subjects were Hispanic/Latino. The mean ages of subjects 6 through 35 months and  
239 36 through 59 months were 21.7 months and 47.1 months, respectively. Subjects in the safety  
240 population (N=2232) received either AFLURIA QUADRIVALENT (N=1673) or a U.S.-  
241 licensed comparator quadrivalent influenza vaccine (N=559). Study subjects were scheduled  
242 to receive either a single vaccination or two vaccinations 28 days apart based on their previous  
243 vaccination history. In this study, AFLURIA QUADRIVALENT and comparator vaccine  
244 were administered by needle and syringe (see *Clinical Studies [14]*).

245 Local (injection site) adverse reactions and systemic adverse events were solicited for 7 days  
246 post-vaccination. Cellulitis-like reactions (defined as concurrent Grade 3 pain, redness, and  
247 swelling/lump) at the injection site were monitored for 28 days post-vaccination. Subjects  
248 were instructed to report and return to clinic within 24 hours in the event of a cellulitis-like  
249 reaction. Unsolicited adverse events were collected for 28 days post-vaccination, and SAEs for  
250 6 months following the last vaccination. All solicited local adverse reactions and systemic  
251 adverse events following any vaccination (first or second dose) are presented in Table 5.

Package insert

252 **Table 5: Proportion of Subjects Per Age Cohort with Any Solicited Local Adverse**  
253 **Reactions or Systemic Adverse Events within 7 Days after Administration of**  
254 **AFLURIA QUADRIVALENT or Comparator QIV (Study 4)<sup>a</sup>**

	Percentage (%) <sup>b</sup> of Subjects in each Age Cohort Reporting an Event							
	6 through 35 months				36 through 59 months			
	AFLURIA Quadrivalent N= 668-669 <sup>c</sup>		Comparator N= 226-227 <sup>c</sup>		AFLURIA Quadrivalent N= 947-949 <sup>c</sup>		Comparator N= 317-318 <sup>c</sup>	
	Any	Gr 3	Any	Gr 3	Any	Gr 3	Any	Gr 3
<b>Local Adverse Reactions</b> <sup>d</sup>								
Pain	20.8	0.1	25.6	0.4	35.5	0	31.4	0.6
Redness	20.8	0.6	17.6	1.8	22.4	2.3	20.8	5.3
Swelling/Lump	6.1	0.4	6.2	0.9	10.1	1.7	12.9	2.5
<b>Systemic Adverse Events</b> <sup>e</sup>								
Irritability	32.9	0.7	28.2	0.4	-	-	-	-
Diarrhea	24.2	0.1	25.6	0.4	12.1	0.1	8.8	0.6
Loss of Appetite	20.0	0.3	19.4	0.4	-	-	-	-
Malaise and Fatigue	-	-	-	-	14.3	0.5	13.2	0.3
Myalgia	-	-	-	-	9.9	0.1	9.4	0
Nausea and/or vomiting	9.4	0.7	11.0	0	9.2	0.4	6.6	0.3
Headache	-	-	-	-	6.2	0.4	5.0	0
Fever <sup>f</sup>	7.2	2.5	11.9	2.6	4.8	1.2	6.0	0.9

255 Abbreviations: Gr 3, Grade 3 (severe); Comparator, Comparator quadrivalent influenza vaccine [Fluzone<sup>®</sup> Quadrivalent  
256 (Sanofi Pasteur)]

257 <sup>a</sup> NCT02914275

258 <sup>b</sup> Percent (%) is derived from the number of subjects that reported the event divided by the number of subjects in the Solicited  
259 Safety Population with non-missing data for each age cohort, treatment group, and each solicited parameter.

260 <sup>c</sup> N = number of subjects in the Solicited Safety Population (subjects who were vaccinated and provided any solicited safety  
261 data) for each study vaccine group.

262 <sup>d</sup> Local adverse reactions: Grade 3 pain is that which prevents daily activity (36 through 59 month subjects); or cried when  
263 limb was moved or spontaneously painful (6 through 35 month subjects); Swelling/Lump and redness: any = ≥ 0mm diameter,  
264 Grade 3 = ≥ 30mm diameter.

265 <sup>e</sup> Systemic adverse events: Fever: any = ≥ 99.5°F (Axillary), Grade 3 = ≥ 101.3°F (Axillary); Grade 3 for all other adverse  
266 events is that which prevents daily activity; Irritability, Loss of Appetite, Malaise and Fatigue, Myalgia and Headache are age  
267 specific systemic adverse events, where “-” denotes event was not applicable to that age cohort.

268 <sup>f</sup> Prophylactic antipyretics (acetaminophen or ibuprophen-containing medications) were not permitted. Antipyretics used to  
269 treat fever were permitted and rates of use were as follows: 6 through 35 months (Afluria QIV 5.9%, Comparator QIV 9.0%);  
270 36 through 59 months (Afluria QIV 3.7%, Comparator QIV 2.5%).

271 In subjects 6 through 35 months of age, all solicited local adverse reactions and systemic  
272 adverse events were reported at lower frequencies after the second vaccination than after the  
273 first vaccination with AFLURIA QUADRIVALENT.

274 In subjects 36 through 59 months of age, all solicited local adverse reactions and systemic  
275 adverse events were reported at lower frequencies after the second vaccination than after the  
276 first vaccination with AFLURIA QUADRIVALENT.

277 The most commonly reported unsolicited adverse events in the 28 days following the first or  
278 second dose of AFLURIA QUADRIVALENT in subjects 6 through 35 months of age were  
279 rhinorrhea (11.2%), cough (10.4%), pyrexia (6.3%), upper respiratory tract infection (4.8%),

**Package insert**

---

280 diarrhea (3.7%), otitis media (2.4%), vomiting (2.4%), nasal congestion (2.4%),  
281 nasopharyngitis (1.9%), irritability (1.7%), ear infection (1.6%), croup infectious (1.4%),  
282 teething (1.3%), rash (1.2%), influenza like illness (1.0%) and fatigue (1.0%), and were similar  
283 to comparator.

284 The most commonly reported unsolicited adverse events in the 28 days following the first or  
285 second dose of AFLURIA QUADRIVALENT in subjects 36 through 59 months of age were  
286 cough (7.7%), rhinorrhea (4.9%), pyrexia (3.7%), upper respiratory tract infection (2.5%),  
287 vomiting (2.1%), nasal congestion (1.6%), nasopharyngitis (1.7%), oropharyngeal pain (1.2%)  
288 diarrhea (1.1%) and fatigue (1.1%), and were similar to the comparator.

289 No deaths were reported in Study 4. In the 180 days following vaccinations, AFLURIA  
290 QUADRIVALENT and comparator vaccine recipients experienced similar rates of serious  
291 adverse events (SAEs), none of which were related to study vaccines. No vaccine-related  
292 febrile seizures occurred in Study 4. Unrelated SAEs of febrile seizures occurred in two  
293 AFLURIA QUADRIVALENT recipients (6 through 35 months age group) at 43 and 104 days  
294 post-vaccinations.

295

**296 6.2 Postmarketing Experience**

297 Because postmarketing reporting of adverse events is voluntary and from a population of  
298 uncertain size, it is not always possible to reliably estimate their frequency or establish a causal  
299 relationship to vaccine exposure. The adverse events described have been included in this  
300 section because they: 1) represent reactions that are known to occur following immunizations  
301 generally or influenza immunizations specifically; 2) are potentially serious; or 3) have been  
302 reported frequently. There are limited postmarketing data available for AFLURIA  
303 QUADRIVALENT. The adverse events listed below reflect experience in both children and  
304 adults and include those identified during post-approval use of AFLURIA (trivalent  
305 formulation) outside the U.S. since 1985.

306 The post-marketing experience with AFLURIA (trivalent formulation) included the following:

**307 Blood and lymphatic system disorders**

308 Thrombocytopenia

**309 Immune system disorders**

310 Allergic or immediate hypersensitivity reactions including anaphylactic shock and serum  
311 sickness

**312 Nervous system disorders**

313 Neuralgia, paresthesia, convulsions (including febrile seizures), encephalomyelitis,  
314 encephalopathy, neuritis or neuropathy, transverse myelitis, and GBS

**315 Vascular disorders**

316 Vasculitis which may be associated with transient renal involvement

**Package insert**

---

317 **Skin and subcutaneous tissue disorders**

318 Pruritus, urticaria, and rash

319 **General disorders and administration site conditions**

320 Cellulitis and large injection site swelling

321 Influenza-like illness

322 **7 DRUG INTERACTIONS**323 No interaction studies have been performed on interaction between influenza vaccines in  
324 general and other vaccines or medications.325 **8 USE IN SPECIFIC POPULATIONS**326 **8.1 Pregnancy**327 Pregnancy Exposure Registry328 There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed  
329 to AFLURIA QUADRIVALENT during pregnancy. Women who are vaccinated with  
330 AFLURIA QUADRIVALENT during pregnancy are encouraged to enroll in the registry by  
331 calling 1-855-358-8966 or sending an email to Seqirus at us.medicalinformation@seqirus.com.  
332333 Risk summary334 All pregnancies have a risk of birth defect, loss, or other adverse outcomes. In the U.S. general  
335 population, the estimated background risk of major birth defects and miscarriage in clinically  
336 recognized pregnancies is 2% to 4% and 15% to 20%, respectively. Data for AFLURIA  
337 (trivalent formulation) administered to pregnant women are relevant to AFLURIA  
338 QUADRIVALENT because both vaccines are manufactured using the same process and have  
339 overlapping compositions (see [Description \[11\]](#)). There are limited data for AFLURIA  
340 QUADRIVALENT administered to pregnant women, and available data for AFLURIA  
341 (trivalent formulation) administered to pregnant women are insufficient to inform vaccine-  
342 associated risks in pregnancy.343 There were no developmental toxicity studies of AFLURIA QUADRIVALENT performed in  
344 animals. A developmental toxicity study of AFLURIA (trivalent formulation) has been  
345 performed in female rats administered a single human dose [0.5 mL (divided)] of AFLURIA  
346 (trivalent formulation) prior to mating and during gestation. This study revealed no evidence  
347 of harm to the fetus due to AFLURIA (trivalent formulation) (see [8.1 Data](#)).348 Clinical Considerations349 *Disease-associated Maternal and/or Embryo-Fetal Risk*350 Pregnant women are at increased risk for severe illness due to influenza compared to non-  
351 pregnant women. Pregnant women with influenza may be at increased risk for adverse  
352 pregnancy outcomes, including preterm labor and delivery.353 Data354 *Animal Data*

**Package insert**

---

355 In a developmental toxicity study, female rats were administered a single human dose [0.5 mL  
356 (divided)] of AFLURIA (trivalent formulation) by intramuscular injection 21 days and 7 days  
357 prior to mating, and on gestation day 6. Some rats were administered an additional dose on  
358 gestation day 20. No vaccine-related fetal malformations or variations and no adverse effects  
359 on pre-weaning development were observed in the study.

**360 8.2 Lactation****361 Risk Summary**

362 It is not known whether AFLURIA QUADRIVALENT is excreted in human milk. Data are  
363 not available to assess the effects of AFLURIA QUADRIVALENT on the breastfed infant or  
364 on milk production/excretion.

365 The developmental and health benefits of breastfeeding should be considered along with the  
366 mother's clinical need for AFLURIA QUADRIVALENT and any potential adverse effects on  
367 the breastfed child from AFLURIA QUADRIVALENT or from the underlying maternal  
368 condition. For preventive vaccines, the underlying maternal condition is susceptibility to  
369 disease prevented by the vaccine.

**370 8.4 Pediatric Use**

371 The safety and effectiveness of AFLURIA QUADRIVALENT in persons less than 6 months of  
372 age have not been established.

373 The PharmaJet Stratis Needle-Free Injection System is not approved as a method of  
374 administering AFLURIA QUADRIVALENT to children and adolescents less than 18 years of  
375 age due to lack of adequate data supporting safety and effectiveness in this population.

**376 8.5 Geriatric Use**

377 In clinical studies, AFLURIA QUADRIVALENT has been administered to, and safety  
378 information collected for, 867 subjects aged 65 years and older (*see Adverse Reactions [6]*).  
379 The 65 years and older age group included 539 subjects 65 through 74 years and 328 subjects  
380 75 years and older. After administration of AFLURIA QUADRIVALENT, hemagglutination-  
381 inhibiting antibody responses were non-inferior to comparator trivalent influenza (TIV-1 and  
382 TIV-2) in persons 65 years of age and older, but were lower than younger adult subjects (*see*  
383 *Clinical Studies [14]*).

384 The PharmaJet Stratis Needle-Free Injection System is not approved as a method of  
385 administering AFLURIA QUADRIVALENT to adults 65 years of age and older due to lack of  
386 adequate data supporting safety and effectiveness in this population.

**387 11 DESCRIPTION**

388 AFLURIA QUADRIVALENT, Influenza Vaccine for intramuscular injection, is a sterile,  
389 clear, colorless to slightly opalescent suspension with some sediment that resuspends upon  
390 shaking to form a homogeneous suspension. AFLURIA QUADRIVALENT is prepared from



**Package insert**

---

391 influenza virus propagated in the allantoic fluid of embryonated chicken eggs. Following  
392 harvest, the virus is purified in a sucrose density gradient using continuous flow zonal  
393 centrifugation. The purified virus is inactivated with beta-propiolactone, and the virus particles  
394 are disrupted using sodium taurodeoxycholate to produce a “split virion”. The disrupted virus  
395 is further purified and suspended in a phosphate buffered isotonic solution.

396 AFLURIA QUADRIVALENT is standardized according to USPHS requirements for the 2018-  
397 2019 influenza season and is formulated to contain 60 mcg hemagglutinin (HA) per 0.5 mL  
398 dose in the recommended ratio of 15 mcg HA for each of the four influenza strains  
399 recommended for the 2018-2019 Northern Hemisphere influenza season:

400 A/Singapore/GP1908/2015 IVR 180A (H1N1) (an A/Michigan/45/2015 – like virus),  
401 A/Singapore/INFIMH-16-0019/2016 IVR-186 (H3N2) (an A/Singapore/INFIMH-16-  
402 0019/2016 – like virus), B/Maryland/15/2016 (a B/Colorado/06/2017 – like virus) and  
403 B/Phuket/3073/2013 BVR-1B (a B/Phuket/3073/2013 – like virus). A 0.25 mL dose contains  
404 7.5 mcg HA of each of the same four influenza strains.

405 Thimerosal, a mercury derivative, is not used in the manufacturing process for the single dose  
406 presentation. This presentation does not contain preservative. The multi-dose presentation  
407 contains thimerosal added as a preservative; each 0.5 mL dose contains 24.5 mcg of mercury  
408 and each 0.25 mL dose contains 12.25 mcg of mercury.

409 A single 0.5 mL dose of AFLURIA QUADRIVALENT contains sodium chloride (4.1 mg),  
410 monobasic sodium phosphate (80 mcg), dibasic sodium phosphate (300 mcg), monobasic  
411 potassium phosphate (20 mcg), potassium chloride (20 mcg), and calcium chloride (0.5 mcg).  
412 From the manufacturing process, each 0.5 mL dose may also contain residual amounts of  
413 sodium taurodeoxycholate ( $\leq 10$  ppm), ovalbumin ( $< 1$  mcg), sucrose ( $< 10$  mcg), neomycin  
414 sulfate ( $\leq 81.8$  nanograms [ng]), polymyxin B ( $\leq 14$  ng), and beta-propiolactone ( $\leq 1.5$  ng). A  
415 single 0.25 mL dose of AFLURIA QUADRIVALENT contains half of these quantities.

416 The rubber tip cap and plunger used for the preservative-free, single-dose syringes and the  
417 rubber stoppers used for the multi-dose vial were not made with natural rubber latex.

**418 12 CLINICAL PHARMACOLOGY****419 12.1 Mechanism of Action**

420 Influenza illness and its complications follow infection with influenza viruses. Global  
421 surveillance of influenza identifies yearly antigenic variants. For example, since 1977  
422 antigenic variants of influenza A (H1N1 and H3N2) and influenza B viruses have been in  
423 global circulation. Since 2001, two distinct lineages of influenza B (Victoria and Yamagata  
424 lineages) have co-circulated worldwide. Specific levels of hemagglutination inhibition (HI)  
425 antibody titers post-vaccination with inactivated influenza vaccine have not been correlated  
426 with protection from influenza virus. In some human studies, antibody titers of 1:40 or greater  
427 have been associated with protection from influenza illness in up to 50% of subjects.<sup>2,3</sup>

**Package insert**

---

428 Antibody against one influenza virus type or subtype confers limited or no protection against  
429 another. Furthermore, antibody to one antigenic variant of influenza virus might not protect  
430 against a new antigenic variant of the same type or subtype. Frequent development of  
431 antigenic variants through antigenic drift is the virologic basis for seasonal epidemics and the  
432 reason for the usual change to one or more new strains in each year's influenza vaccine.  
433 Therefore, inactivated influenza vaccines are standardized to contain the HA of four strains  
434 (i.e., typically two type A and two type B) representing the influenza viruses likely to be  
435 circulating in the U.S. during the upcoming winter.

436 Annual revaccination with the current vaccine is recommended because immunity declines  
437 during the year after vaccination and circulating strains of influenza virus change from year to  
438 year.<sup>1</sup>

**439 13 NONCLINICAL TOXICOLOGY****440 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

441 AFLURIA QUADRIVALENT has not been evaluated for carcinogenic or mutagenic potential,  
442 or male infertility in animals. A developmental toxicity study conducted in rats vaccinated  
443 with AFLURIA (trivalent formulation) revealed no impact on female fertility (see [Pregnancy](#)  
444 [\[8.1\]](#)).

**445 14 CLINICAL STUDIES****446 14.1 Efficacy Against Laboratory-Confirmed Influenza**

447 The efficacy of AFLURIA (trivalent formulation) is relevant to AFLURIA QUADRIVALENT  
448 because both vaccines are manufactured using the same process and have overlapping  
449 compositions (see [Description \[11\]](#)).

450 The efficacy of AFLURIA (trivalent formulation) was demonstrated in Study 5, a randomized,  
451 observer-blind, placebo-controlled study conducted in 15,044 subjects. Healthy subjects 18  
452 through 64 years of age were randomized in a 2:1 ratio to receive a single dose of AFLURIA  
453 (trivalent formulation) (enrolled subjects: 10,033; evaluable subjects: 9,889) or placebo  
454 (enrolled subjects: 5,011; evaluable subjects: 4,960). The mean age of all randomized subjects  
455 was 35.5 years. 54.4% were female and 90.2% were White. Laboratory-confirmed influenza  
456 was assessed by active and passive surveillance of influenza-like illness (ILI) beginning 2  
457 weeks post-vaccination until the end of the influenza season, approximately 6 months post-  
458 vaccination. ILI was defined as at least one respiratory symptom (e.g., cough, sore throat,  
459 nasal congestion) and at least one systemic symptom (e.g., oral temperature of 100.0°F or  
460 higher, feverishness, chills, body aches). Nasal and throat swabs were collected from subjects  
461 who presented with an ILI for laboratory confirmation by viral culture and real-time reverse  
462 transcription polymerase chain reaction. Influenza virus strain was further characterized using  
463 gene sequencing and pyrosequencing.

464 Attack rates and vaccine efficacy, defined as the relative reduction in the influenza infection  
465 rate for AFLURIA (trivalent formulation) compared to placebo, were calculated using the per

**Package insert**

466 protocol population. Vaccine efficacy against laboratory-confirmed influenza infection due to  
467 influenza A or B virus strains contained in the vaccine was 60% with a lower limit of the 95%  
468 CI of 41% (Table 6).

469 **Table 6: AFLURIA (trivalent formulation): Laboratory-Confirmed Influenza Infection**  
470 **Rate and Vaccine Efficacy in Adults 18 through 64 Years of Age (Study 5)<sup>a</sup>**

	Subjects <sup>b</sup>	Laboratory-Confirmed Influenza Cases	Influenza Infection Rate	Vaccine Efficacy <sup>c</sup>	
	N			N	n/N %
<b>Vaccine-matched Strains</b>					
AFLURIA	9889	58	0.59	60	41
Placebo	4960	73	1.47		
<b>Any Influenza Virus Strain</b>					
AFLURIA	9889	222	2.24	42	28
Placebo	4960	192	3.87		

471 Abbreviations: CI, confidence interval.

472 <sup>a</sup> NCT00562484

473 <sup>b</sup> The Per Protocol Population was identical to the Evaluable Population in this study.

474 <sup>c</sup> Vaccine efficacy = 1 minus the ratio of AFLURIA (trivalent formulation) /placebo infection rates. The objective of the study  
475 was to demonstrate that the lower limit of the CI for vaccine efficacy was greater than 40%.

476 **14.2 Immunogenicity of AFLURIA QUADRIVALENT in Adults and Older Adults**  
477 **Administered by Needle and Syringe**

478 Study 1 was a randomized, double-blind, active-controlled trial conducted in the U.S. in adults  
479 aged 18 years of age and older. Subjects received one dose of either AFLURIA  
480 QUADRIVALENT (N=1691) or one of two formulations of comparator trivalent influenza  
481 vaccine (AFLURIA, TIV-1 N=854 or TIV-2 N=850) each containing an influenza type B virus  
482 that corresponded to one of the two B viruses in AFLURIA QUADRIVALENT (a type B virus  
483 of the Yamagata lineage or a type B virus of the Victoria lineage, respectively).

484 Post-vaccination immunogenicity was evaluated on sera obtained 21 days after administration  
485 of a single dose of AFLURIA QUADRIVALENT or TIV comparator. The co-primary  
486 endpoints were HI Geometric Mean Titer (GMT) ratios (adjusted for baseline HI titers) and the  
487 difference in seroconversion rates for each vaccine strain, 21 days after vaccination. Pre-  
488 specified non-inferiority criteria required that the upper bound of the 2-sided 95% CI of the  
489 GMT ratio (TIV/AFLURIA QUADRIVALENT) did not exceed 1.5 and the upper bound of the  
490 2-sided 95% CI of the seroconversion rate difference (TIV minus AFLURIA  
491 QUADRIVALENT) did not exceed 10.0% for each strain.

492 Serum HI antibody responses to AFLURIA QUADRIVALENT were non-inferior to both TIVs  
493 for all influenza strains for subjects 18 years of age and older. Additionally, non-inferiority  
494 was demonstrated for both endpoints in both age sub-groups, adults aged 18 through 64 years  
495 and 65 years and older, for all strains (Table 7). Superiority of the immune response to each of  
496 the influenza B strains contained in AFLURIA QUADRIVALENT was shown relative to the

**Package insert**

---

497 antibody response after vaccination with TIV formulations not containing that B lineage strain  
498 for subjects 18 years of age and older. Superiority against the alternate B strain was also  
499 demonstrated for each of the influenza B strains in both age sub-groups; 18 through 64 years  
500 and 65 years and older. Post-hoc analyses of immunogenicity endpoints by gender did not  
501 demonstrate meaningful differences between males and females. The study population was not  
502 sufficiently diverse to assess differences between races or ethnicities.

Package insert

503 **Table 7: Post-Vaccination HI Antibody GMTs, Seroconversion Rates, and Analyses of**  
504 **Non-Inferiority of AFLURIA QUADRIVALENT Relative to Trivalent**  
505 **Influenza Vaccine (TIV) by Age Cohort (Study 1)<sup>a</sup>**

Strain	Post-vaccination GMT		GMT Ratio <sup>b</sup>	Seroconversion % <sup>c</sup>		Difference	Met both pre-defined non-inferiority criteria? <sup>d</sup>
	AFLURIA Quadrivalent	Pooled TIV or TIV-1 (B Yamagata) or TIV-2 (B Victoria)	Pooled TIV or TIV-1 or TIV-2 over AFLURIA Quadrivalent (95% CI)	AFLURIA Quadrivalent N=1691	Pooled TIV or TIV-1 (B Yamagata) or TIV-2 (B Victoria)	Pooled TIV or TIV-1 or TIV-2 minus AFLURIA Quadrivalent (95% CI)	
<b>18 through 64 years</b>	<b>AFLURIA Quadrivalent N=835, Pooled TIV N=845, TIV-1 N=424, TIV-2 N=421</b>						
A(H1N1)	432.7	402.8	0.93 <sup>e</sup> (0.85, 1.02)	51.3	49.1	-2.1 <sup>h</sup> (-6.9, 2.7)	Yes
A(H3N2)	569.1	515.1	0.91 <sup>e</sup> (0.83, 0.99)	56.3	51.7	-4.6 <sup>h</sup> (-9.4, 0.2)	Yes
B/Massachusetts/2/2012 (B Yamagata)	92.3	79.3	0.86 <sup>f</sup> (0.76, 0.97)	45.7	41.3	-4.5 <sup>i</sup> (-10.3, 1.4)	Yes
B/Brisbane/60/2008 (B Victoria)	110.7	95.2	0.86 <sup>g</sup> (0.76, 0.98)	57.6	53.0	-4.6 <sup>j</sup> (-10.5, 1.2)	Yes
<b>≥ 65 years</b>	<b>AFLURIA Quadrivalent N=856, Pooled TIV N=859, TIV-1 N=430, TIV-2 N=429</b>						
A(H1N1)	211.4	199.8	0.95 <sup>e</sup> (0.88, 1.02)	26.6	26.4	-0.2 <sup>h</sup> (-5.0, 4.5)	Yes
A(H3N2)	419.5	400.0	0.95 <sup>e</sup> (0.89, 1.02)	25.9	27.0	1.1 <sup>h</sup> (-3.7, 5.8)	Yes
B/Massachusetts/2/2012 (B Yamagata)	43.3	39.1	0.90 <sup>f</sup> (0.84, 0.97)	16.6	14.4	-2.2 <sup>i</sup> (-8.0, 3.6)	Yes
B/Brisbane/60/2008 (B Victoria)	66.1	68.4	1.03 <sup>g</sup> (0.94, 1.14)	23.5	24.7	1.2 <sup>j</sup> (-4.6, 7.0)	Yes

506 Abbreviations: CI, confidence interval; GMT, geometric mean titer.

507 <sup>a</sup> NCT02214225

508 <sup>b</sup> GMT ratio was computed after fitting a multi-variable model on the post-vaccination titers including sex, vaccination history,  
509 pre-vaccination HI titers and other factors.

510 <sup>c</sup> Seroconversion rate is defined as a 4-fold increase in post-vaccination HI antibody titer from pre-vaccination titer  $\geq 1:10$  or an  
511 increase in titer from  $< 1:10$  to  $\geq 1:40$ .

512 <sup>d</sup> Non-inferiority (NI) criterion for the GMT ratio: upper bound of 2-sided 95% CI on the GMT ratio of Pooled TIV or TIV-1 (B  
513 Yamagata) or TIV-2 (B Victoria)/AFLURIA Quadrivalent should not exceed 1.5. NI criterion for the SCR difference: upper  
514 bound of 2-sided 95% CI on the difference between SCR Pooled TIV or TIV-1 (B Yamagata) or TIV-2 (B Victoria) minus  
515 AFLURIA Quadrivalent should not exceed 10%.

516 <sup>e</sup> Pooled TIV/AFLURIA Quadrivalent

517 <sup>f</sup> TIV-1 (B Yamagata)/AFLURIA Quadrivalent

518 <sup>g</sup> TIV-2 (B Victoria)/AFLURIA Quadrivalent

519 <sup>h</sup> Pooled TIV – AFLURIA Quadrivalent

520 <sup>i</sup> TIV-1 (B Yamagata) - AFLURIA Quadrivalent

521 <sup>j</sup> TIV-2 (B Victoria) - AFLURIA Quadrivalent

Package insert

**14.3 Immunogenicity of AFLURIA (trivalent formulation) Administered by PharmaJet Stratis Needle-Free Injection System**

Study 2 was a randomized, comparator-controlled, non-inferiority study that enrolled 1,250 subjects 18 through 64 years of age. This study compared the immune response following administration of AFLURIA (trivalent formulation) when delivered intramuscularly using either the PharmaJet Stratis Needle-Free Injection System or needle and syringe. Immunogenicity assessments were performed prior to vaccination and at 28 days after vaccination in the immunogenicity population (1130 subjects, 562 PharmaJet Stratis Needle-Free Injection System group, 568 needle and syringe group). The co-primary endpoints were HI GMT ratios for each vaccine strain and the absolute difference in seroconversion rates for each vaccine strain 28 days after vaccination. As shown in Table 8, non-inferiority of administration of AFLURIA (trivalent formulation) by the PharmaJet Stratis Needle-Free Injection System compared to administration of AFLURIA (trivalent formulation) by needle and syringe was demonstrated in the immunogenicity population for all strains. Post-hoc analyses of immunogenicity by age showed that younger subjects (18 through 49 years) elicited higher immunological responses than older subjects (50 through 64 years). Post-hoc analyses of immunogenicity according to sex and body mass index did not reveal significant influences of these variables on immune responses. The study population was not sufficiently diverse to assess immunogenicity by race or ethnicity.

**Table 8: Baseline and Post-Vaccination HI Antibody GMTs, Seroconversion Rates, and Analyses of Non-Inferiority of AFLURIA (trivalent formulation) Administered by PharmaJet Stratis Needle-Free Injection System or Needle and Syringe, Adults 18 through 64 Years of Age (Study 2)<sup>a</sup>**

Strain	Baseline GMT		Post-vaccination GMT		GMT Ratio <sup>b</sup>	Seroconversion % <sup>c</sup>		Difference	Met both pre-defined non-inferiority criteria? <sup>d</sup>
	Needle and Syringe N=568	PharmaJet Stratis Needle-Free Injection System N=562	Needle and Syringe N=568	PharmaJet Stratis Needle-Free Injection System N=562	Needle and Syringe over PharmaJet Stratis Needle-Free Injection System (95% CI)	Needle and Syringe N=568	PharmaJet Stratis Needle-Free Injection System N=562	Needle and Syringe minus PharmaJet Stratis Needle-Free Injection System (95% CI)	
A(H1N1)	79.5	83.7	280.6	282.9	0.99 (0.88, 1.12)	38.4	37.5	0.8 (-4.8, 6.5)	Yes
A(H3N2)	75.4	68.1	265.9	247.3	1.08 (0.96, 1.21)	45.1	43.8	1.3 (-4.5, 7.1)	Yes
B	12.6	13.5	39.7	42.5	0.94 (0.83, 1.06)	35.2	34.9	0.3 (-5.2, 5.9)	Yes

Abbreviations: CI, confidence interval; GMT, geometric mean titer.

<sup>a</sup> NCT01688921

<sup>b</sup> GMT ratio is defined as post-vaccination GMT for Needle and Syringe/PharmaJet Stratis Needle-Free Injection System.

<sup>c</sup> Seroconversion rate is defined as a 4-fold increase in post-vaccination HI antibody titer from pre-vaccination titer  $\geq 1:10$  or an increase in titer from  $< 1:10$  to  $\geq 1:40$ .

<sup>d</sup> Non-inferiority (NI) criterion for the GMT ratio: upper bound of 2-sided 95% CI on the GMT ratio of Needle and Syringe/PharmaJet Stratis Needle-Free Injection System should not exceed 1.5. NI criterion for the seroconversion rate (SCR) difference: upper bound of 2-sided 95% CI on the difference between SCR Needle and Syringe – SCR PharmaJet

**Package insert**

---

553 Stratis Needle-Free Injection System should not exceed 10%.

554 **14.4 Immunogenicity of AFLURIA QUADRIVALENT in Children 5 through 17**  
555 **Years Administered by Needle and Syringe**

556 Study 3 was a randomized, observer-blinded, comparator-controlled trial conducted in the U.S.  
557 in children 5 through 17 years of age. A total of 2278 subjects were randomized 3:1 to receive  
558 one or two doses of AFLURIA QUADRIVALENT (N=1709) or a U.S.-licensed comparator  
559 quadrivalent influenza vaccine (N=569). Subjects 5 through 8 years of age were eligible to  
560 receive a second dose at least 28 days after the first dose depending on their influenza  
561 vaccination history, consistent with the 2015-2016 recommendations of the Advisory  
562 Committee on Immunization Practices (ACIP) for Prevention and Control of Seasonal  
563 Influenza with Vaccines. Approximately 25% of subjects in each treatment group in the 5  
564 through 8 years of age sub-group received two vaccine doses.

565 Baseline serology for HI assessment was collected prior to vaccination. Post-vaccination  
566 immunogenicity was evaluated by HI assay on sera obtained 28 days after the last vaccination  
567 dose.

568 The primary objective was to demonstrate that vaccination with AFLURIA QUADRIVALENT  
569 elicits an immune response that is not inferior to that of a comparator vaccine containing the  
570 same recommended virus strains. The Per Protocol Population (AFLURIA QUADRIVALENT  
571 n=1605, Comparator n=528) was used for the primary endpoint analyses. The co-primary  
572 endpoints were HI Geometric Mean Titer (GMT) ratios (adjusted for baseline HI titers and  
573 other covariates) and seroconversion rates for each vaccine strain, 28 days after the last  
574 vaccination. Pre-specified non-inferiority criteria required that the upper bound of the 2-sided  
575 95% CI of the GMT ratio (Comparator/AFLURIA QUADRIVALENT) did not exceed 1.5 and  
576 the upper bound of the 2-sided 95% CI of the seroconversion rate difference (Comparator  
577 minus AFLURIA QUADRIVALENT) did not exceed 10.0% for each strain. Serum HI  
578 antibody responses to AFLURIA QUADRIVALENT were non-inferior for both GMT ratio  
579 and seroconversion rates relative to the comparator vaccine for all influenza strains (Table 9).  
580 Analyses of immunogenicity endpoints by gender did not demonstrate meaningful differences  
581 between males and females. The study population was not sufficiently diverse to assess  
582 differences among races or ethnicities.

Package insert

583 **Table 9: Post-Vaccination HI Antibody GMTs, SCRs, and Analyses of Non-Inferiority of**  
584 **AFLURIA QUADRIVALENT Relative to a U.S.-Licensed Comparator**  
585 **Quadrivalent Influenza Vaccine for each Strain 28 Days after Last Vaccination**  
586 **Among a Pediatric Population 5 through 17 Years of Age (Per Protocol**  
587 **Population) (Study 3) <sup>a,b</sup>**

Strain	Post-vaccination GMT		GMT Ratio <sup>c</sup>	Seroconversion % <sup>d</sup>		SCR Difference <sup>e</sup>	Met both pre-defined non-inferiority criteria? <sup>f</sup>
	AFLURIA Quadrivalent N=1605	Comparator N=528	Comparator over AFLURIA Quadrivalent (95% CI)	AFLURIA Quadrivalent N=1605 (95% CI)	Comparator N=528 (95% CI)	Comparator minus AFLURIA Quadrivalent (95% CI)	
A(H1N1)	952.6 (n=1604 <sup>g</sup> )	958.8	1.01 (0.93, 1.09)	66.4 (64.0, 68.7)	63.3 (59.0, 67.4)	-3.1 (-8.0, 1.8)	Yes
A(H3N2)	886.4 (n=1604 <sup>g</sup> )	930.6	1.05 (0.96, 1.15)	82.9 (81.0, 84.7)	83.3 (79.9, 86.4)	0.4 (-4.5, 5.3)	Yes
B/Phuket/3073/2013 (B Yamagata)	60.9 (n=1604 <sup>g</sup> )	54.3	0.89 (0.81, 0.98)	58.5 (56.0, 60.9)	55.1 (50.8, 59.4)	-3.4 (-8.3, 1.5)	Yes
B/Brisbane/60/2008 (B Victoria)	145.0 (n=1604 <sup>g</sup> )	133.4	0.92 (0.83, 1.02)	72.1 (69.8, 74.3)	70.1 (66.0, 74.0)	-2.0 (-6.9, 2.9)	Yes

588 Abbreviations: CI, confidence interval; Comparator, Comparator quadrivalent influenza vaccine (Fluarix<sup>®</sup> Quadrivalent  
589 [GlaxoSmithKline Biologicals]); GMT (adjusted), geometric mean titer; SCR, seroconversion rate.

590 <sup>a</sup> NCT02545543

591 <sup>b</sup> The Per-Protocol Population comprised all subjects in the Evaluable Population who did not have any protocol deviations  
592 that were medically assessed as potentially impacting on immunogenicity results.

593 <sup>c</sup> GMT Ratio = Comparator /AFLURIA QUADRIVALENT. Adjusted analysis model: Log-transformed Post-Vaccination HI  
594 Titer=Vaccine + Age Strata [5-8, 9-17] + Gender + Vaccination History [y/n] + Log-transformed Pre-Vaccination HI Titer +  
595 Site + Number of Doses (1 vs 2) + Age Strata\*Vaccine. The Age Strata\*Vaccine interaction term was excluded from the  
596 model fit for the strains B/Yamagata and B/Victoria as the interaction result was non-significant (p>0.05). Least square  
597 means were back transformed.

598 <sup>d</sup> Seroconversion rate was defined as the percentage of subjects with either a prevaccination HI titer < 1:10 and a  
599 postvaccination HI titer ≥ 1:40 or a prevaccination HI titer ≥ 1:10 and a 4-fold increase in postvaccination HI titer.

600 <sup>e</sup> Seroconversion rate difference = Comparator SCR percentage minus AFLURIA QUADRIVALENT SCR percentage.

601 <sup>f</sup> Non-inferiority (NI) criterion for the GMT ratio: upper bound of two-sided 95% CI on the GMT ratio of Comparator  
602 /AFLURIA QUADRIVALENT should not exceed 1.5. NI criterion for the SCR difference: upper bound of two-sided 95%  
603 CI on the difference between SCR Comparator – AFLURIA QUADRIVALENT should not exceed 10%.

604 <sup>g</sup> Subject 8400394-0046 was excluded from the Per-Protocol Population for the adjusted GMT analysis for the GMT ratio since  
605 the subject did not have information on all covariates (unknown prevaccination history).

606 **14.5 Immunogenicity of AFLURIA QUADRIVALENT in Children 6 Months**  
607 **through 59 Months Administered by Needle and Syringe**

608 Study 4 was a randomized, observer-blind, comparator-controlled trial conducted in the U.S.  
609 in children 6 months through 59 months of age. A total of 2247 subjects were randomized 3:1  
610 to receive AFLURIA QUADRIVALENT (N=1684) or a U.S.-licensed comparator  
611 quadrivalent influenza vaccine (N=563). Children 6 months through 35 months received one  
612 or two 0.25 mL doses and children 36 months through 59 months received one or two 0.5 mL  
613 doses. Subjects were eligible to receive a second dose at least 28 days after the first dose  
614 depending on their influenza vaccination history, consistent with the 2016-2017  
615 recommendations of the Advisory Committee on Immunization Practices (ACIP) for



**Package insert**

---

616 Prevention and Control of Seasonal Influenza with Vaccines. Approximately 40% of subjects  
617 in each treatment group received two vaccine doses.

618 Baseline serology for HI assessment was collected prior to vaccination. Postvaccination  
619 immunogenicity was evaluated by HI assay on sera obtained 28 days after the last vaccination  
620 dose.

621 The primary objective was to demonstrate that vaccination with AFLURIA  
622 QUADRIVALENT elicits an immune response that is not inferior to that of a comparator  
623 vaccine containing the same recommended virus strains. The Per Protocol Population  
624 (AFLURIA QUADRIVALENT n=1456, Comparator QIV n=484) was used for the primary  
625 endpoint analyses. The co-primary endpoints were HI Geometric Mean Titer (GMT) ratios  
626 (adjusted for baseline HI titers and other covariates) and seroconversion rates for each vaccine  
627 strain, 28 days after the last vaccination. Pre-specified non-inferiority criteria required that  
628 the upper bound of the 2-sided 95% CI of the GMT ratio (Comparator QIV/AFLURIA  
629 QUADRIVALENT) did not exceed 1.5 and the upper bound of the 2-sided 95% CI of the  
630 seroconversion rate difference (Comparator QIV minus AFLURIA QUADRIVALENT) did  
631 not exceed 10.0% for each strain. Serum HI antibody responses to AFLURIA  
632 QUADRIVALENT were non-inferior for both GMT ratio and seroconversion rates relative to  
633 the comparator vaccine for all influenza strains (Table 10). Analyses of immunogenicity  
634 endpoints by gender did not demonstrate meaningful differences between males and females.  
635 The study population was not sufficiently diverse to assess differences among races or  
636 ethnicities.

Package insert

637 **Table 10: Post-Vaccination HI Antibody GMTs, SCR, and Analyses of Non-Inferiority**  
 638 **of AFLURIA QUADRIVALENT Relative to a U.S.-Licensed Comparator**  
 639 **Quadrivalent Influenza Vaccine for each Strain 28 Days after Last**  
 640 **Vaccination Among a Pediatric Population 6 through 59 Months of Age (Per**  
 641 **Protocol Population) (Study 4)<sup>a, b</sup>**

Strain	Post-vaccination GMT		GMT Ratio <sup>c</sup>	Seroconversion % <sup>d</sup>		SCR Difference <sup>e</sup>	Met both pre-defined non-inferiority criteria? <sup>f</sup>
	AFLURIA Quadrivalent N=1456	Comparator N=484	Comparator over AFLURIA Quadrivalent (95% CI)	AFLURIA Quadrivalent N=1456 (95% CI)	Comparator N=484 (95% CI)	Comparator minus AFLURIA Quadrivalent (95% CI)	
A(H1N1)	353.5 (n=1455 <sup>g</sup> )	281.0 (n=484)	0.79 (0.72, 0.88)	79.1 (76.9, 81.1) (n=1456)	68.8 (64.5, 72.9) (n=484)	-10.3 (-15.4, -5.1)	Yes
A(H3N2)	393.0 (n=1454 <sup>g</sup> )	500.5 (n=484)	1.27 (1.15, 1.42)	82.3 (80.2, 84.2) (n=1455 <sup>h</sup> )	84.9 (81.4, 88.0) (n=484)	2.6 (-2.5, 7.8)	Yes
B/Phuket/3073/2013 (B Yamagata)	23.7 (n=1455 <sup>g</sup> )	26.5 (n=484)	1.12 (1.01, 1.24)	38.9 (36.4, 41.4) (n=1456)	41.9 (37.5, 46.5) (n=484)	3.1 (-2.1, 8.2)	Yes
B/Brisbane/60/2008 (B Victoria)	54.6 (n=1455 <sup>g</sup> )	52.9 (n=483 <sup>h</sup> )	0.97 (0.86, 1.09)	60.2 (57.6, 62.7) (n=1456)	61.1 (56.6, 65.4) (n=483 <sup>h</sup> )	0.9 (-4.2, 6.1)	Yes

642 Abbreviations: CI, confidence interval; Comparator, Comparator quadrivalent influenza vaccine (Fluzone Quadrivalent  
 643 [Sanofi Aventis]); GMT (adjusted), geometric mean titer; SCR, seroconversion rate.

644 <sup>a</sup> NCT02914275

645 <sup>b</sup> The Per-Protocol Population comprised all subjects (6 through 35 months of age receiving one or two 0.25 mL doses and 36  
 646 through 59 months of age receiving one or two 0.5 mL doses) in the Evaluable Population who did not have any protocol  
 647 deviations that were medically assessed as potentially impacting on immunogenicity results.

648 <sup>c</sup> GMT Ratio = Comparator / AFLURIA QUADRIVALENT. Adjusted analysis model: Log-transformed Post-Vaccination HI  
 649 Titer=Vaccine + Age Cohort [6 through 35 months or 36 through 59 months] + Gender + Vaccination History [y/n] + Log-  
 650 transformed Pre-Vaccination HI Titer + Site + Number of Doses (1 vs 2) + Age Cohort\*Vaccine. The Age Cohort\*Vaccine  
 651 interaction term was excluded from the model fit for the strains A(H1N1), A(H3N2) and B/Yamagata as the interaction result  
 652 was non-significant (p>0.05). Least square means were back transformed.

653 <sup>d</sup> Seroconversion rate was defined as the percentage of subjects with either a prevaccination HI titer < 1:10 and a  
 654 postvaccination HI titer ≥ 1:40 or a prevaccination HI titer ≥ 1:10 and a 4-fold increase in postvaccination HI titer.

655 <sup>e</sup> Seroconversion rate difference = Comparator SCR percentage minus AFLURIA QUADRIVALENT SCR percentage.

656 <sup>f</sup> Noninferiority (NI) criterion for the GMT ratio: upper bound of two-sided 95% CI on the GMT ratio of Comparator /  
 657 AFLURIA QUADRIVALENT should not exceed 1.5. NI criterion for the SCR difference: upper bound of two-sided 95% CI  
 658 on the difference between SCR Comparator – AFLURIA QUADRIVALENT should not exceed 10%.

659 <sup>g</sup> Subject 8400402-0073 was excluded from the Per-Protocol Population for the adjusted GMT analysis for the GMT ratio  
 660 because the subject did not have information on all covariates (unknown prevaccination history).

661 <sup>h</sup> Subject 8400427-0070 had missing B/Victoria Antigen pre-vaccination titer.

662 <sup>i</sup> Subject 8400402-0074 had missing A/H3N2 post-vaccination titer.

663 **15 REFERENCES**

- 664 1. Centers for Disease Control and Prevention. Prevention and Control of Influenza:  
 665 Recommendations of the Advisory Committee on Immunization Practices (ACIP).  
 666 *MMWR Recomm Rep* 2010;59 (RR-8):1-62.  
 667 2. Hannoun C, Megas F, Piercy J. Immunogenicity and Protective Efficacy of Influenza  
 668 Vaccination. *Virus Res* 2004;103:133-138.  
 669 3. Hobson D, Curry RL, Beare AS, et al. The Role of Serum Hemagglutination-  
 670 Inhibiting Antibody in Protection against Challenge Infection with Influenza A2 and B  
 671 Viruses. *J Hyg Camb* 1972;70:767-777.

**Package insert**672 **16 HOW SUPPLIED/STORAGE AND HANDLING**673 **16.1 How Supplied**

674 Each product presentation includes a package insert and the following components:

Presentation	Carton NDC Number	Components
Pre-Filled Syringe	33332-218-20	<ul style="list-style-type: none"><li>Ten 0.25 mL single-dose syringes fitted with a Luer-Lok™ attachment without needles [NDC 33332-218-21]</li></ul>
Pre-Filled Syringe	33332-318-01	<ul style="list-style-type: none"><li>Ten 0.5 mL single-dose syringes fitted with a Luer-Lok™ attachment without needles [NDC 33332-318-02]</li></ul>
Multi-Dose Vial	33332-418-10	<ul style="list-style-type: none"><li>One 5 mL vial, which contains ten 0.5 mL doses [NDC 33332-418-11]</li></ul>

675 **16.2 Storage and Handling**

- 676 • Store refrigerated at 2–8°C (36–46°F).
- 677 • Do not freeze. Discard if product has been frozen.
- 678 • Protect from light.
- 679 • Do not use AFLURIA QUADRIVALENT beyond the expiration date printed on the
- 680 label.
- 681 • Between uses, return the multi-dose vial to the recommended storage conditions.
- 682 • Once the stopper of the multi-dose vial has been pierced the vial must be discarded
- 683 within 28 days.

684 **17 PATIENT COUNSELING INFORMATION**

- 685 • Inform the vaccine recipient or guardian of the potential benefits and risks of
- 686 immunization with AFLURIA QUADRIVALENT.
- 687 • Inform the vaccine recipient or guardian that AFLURIA QUADRIVALENT is an
- 688 inactivated vaccine that cannot cause influenza but stimulates the immune system to
- 689 produce antibodies that protect against influenza, and that the full effect of the vaccine
- 690 is generally achieved approximately 3 weeks after vaccination.
- 691 • Instruct the vaccine recipient or guardian to report any severe or unusual adverse
- 692 reactions to their healthcare provider.
- 693 • Encourage women who receive AFLURIA QUADRIVALENT while pregnant to
- 694 enroll in the pregnancy registry. Pregnant women can enroll in the pregnancy registry
- 695 by calling 1-855-358-8966 or sending an email to Seqirus at
- 696 us.medicalinformation@seqirus.com.
- 697 • Provide the vaccine recipient Vaccine Information Statements prior to immunization.
- 698 These materials are available free of charge at the Centers for Disease Control and
- 699 Prevention (CDC) website ([www.cdc.gov/vaccines](http://www.cdc.gov/vaccines)).
- 700 • Instruct the vaccine recipient that annual revaccination is recommended.



**Package insert**

---

701 Manufactured by:  
702 **Seqirus Pty Ltd.** Parkville, Victoria, 3052, Australia  
703 U.S. License No. 2044

704 Distributed by:  
705 **Seqirus USA Inc.** 25 Deforest Avenue, Summit, NJ 07901, USA  
706 1-855-358-8966

707 AFLURIA is a registered trademark of Seqirus UK Limited or its affiliates.  
708 PharmaJet® and STRATIS® are registered trademarks of PharmaJet.  
709 Luer-Lok™ is a trademark of Becton, Dickinson and Company Corporation.